



Further advice on Scarlett fever

Dear Parent/Career

We are writing in response to the rising cases of Scarlet fever. Although scarlet fever is usually a mild illness, it should be treated with antibiotics to minimise the risk of complications and reduce the spread to others.

The symptoms of scarlet fever include a sore throat, headache, fever, nausea and vomiting. This is followed by a fine red rash which typically first appears on the chest and stomach, rapidly spreading to other parts of the body. On more darkly pigmented skin, the scarlet rash may be harder to spot, but it should feel like 'sandpaper'. The face can be flushed red but pale around the mouth.

If you think you, or your child, have scarlet fever:

- see your GP or contact NHS 111 as soon as possible.
- make sure that you/your child takes the full course of any antibiotics prescribed by the doctor.
- stay at home, away from nursery, school or work for at least 24 hours after starting the antibiotic treatment, to avoid spreading the infection.

Children who have had chickenpox recently are more likely to develop more serious infection during an outbreak of scarlet fever and so parents/carers should remain vigilant for symptoms such as a persistent high fever, cellulitis (skin infection) and arthritis (joint pain and swelling). If you are concerned for any reason please seek medical assistance immediately. If your child has an underlying condition which affects their immune system, you should contact your GP or hospital doctor to discuss whether any additional measures are needed.

Cases that arise in school are reported to UKHSA and affected areas are then steam cleaned.

Yours sincerely

Mr. K Baffoe

BJS FEDERATION OF SCHOOLS

Bonneville, Jessop & Stockwell Primary Schools

Jessop Primary School
Lowden Road
London SE24 0BJ

Tel: 02072742333
Email: admin@jessop.lambeth.sch.uk
Web: jessop.lambeth.sch.uk

Executive Headteacher
Andrea Parker, BA (Hons), NPQH, NPQEL
Head of School
Kenneth Baffoe